

The Effect of Sodium Nitrite Exposure on Physiological Response of Starved Far Eastern Catfish, *Silurus asotus*

†In-Seok Park¹ and Hyun Woo Gil²

¹Division of Marine Bioscience, College of Ocean Science and Technology, Korea Maritime and Ocean University, Busan 49112, Korea

²Bio-Monitoring Center, Sejong 30121, Korea

ABSTRACT : The experiment was conducted for 210 days to determine the effect of feeding, and starvation, and exposure to sodium nitrite (NaNO₂) on the survival, physiological changes, hematological parameter, and stress response of Far Eastern catfish, *Silurus asotus*. The survival of the starved group was lower than that of the fed group during the experiment. Starvation resulted in retardation of growth, which provides an example of fish that failed to continue to grow and remain in a good condition. Blood analyses (cortisol and glucose) showed significant differences of stress response between the fed and starved groups exposed to NaNO₂ at the conclusion of the experiment ($p < 0.05$). These results suggest that all nutritional parameters used for starvation and feeding with NaNO₂ stress in this experiment appear to be a useful index of nutritional status in Far Eastern catfish.

Key words : Far Eastern catfish (*Silurus asotus*), Physiological response, Sodium nitrite, Starvation

INTRODUCTION

Most fish undergo the period of starvation because of wintering, spawning, migration, or regional decrease in food resources (Mustafa & Mittal, 1982; Park et al., 2001, 2015). Fish can overcome starvation through biochemical, physiological, and behavioral strategies. Endogenous energy from basic metabolic accumulation in the body is spent as fish consume their own tissues to remain alive during starvation (Mustafa & Mittal, 1982; Weatherley & Gill, 1987; Kim et al., 2012; Park et al., 2013). During starvation, essential processes in fish are maintained at the expense of accumulated (i.e., completely endogenous) energy reserves, resulting in the progressive depletion and wast-

age (degrowth) of body tissues (Weatherley & Gill, 1987; Park et al., 2015). The observed incidence of starvation is fundamentally common based on morphological and histological criteria (Park et al., 2001).

The efficiency of oxygen consumption in fish is directly related to their metabolic processes, through which they synthesize body ingredients to produce physical energy. This, in turn, ultimately determines their population density, food supply, and fish yield (Mehner & Wieser, 1994). Mehner and Wieser (1994) observed the relationship between the depletion of energy reserves and changes in oxygen consumption in the perch, *Perca fluviatilis*. Starvation reduces oxygen consumption efficiency as reported in perch larvae and the traira, *Hoplias malabaricus*, when

Manuscript received September 28, 2018, Received in revised form October 15, 2018, Accepted November 2, 2018

† Corresponding Author : In-Seok Park, Division of Marine Bioscience, College of Ocean Science and Technology, Korea Maritime and Ocean University, Busan 49112, Korea. Tel: +82- 51-410-4321, Fax: +82-51-404-4750, E-mail: ispark@kmou.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

they were subjected to extended periods of starvation (Mehner & Wieser, 1994; Rios et al., 2002).

Stresses on fish induce the release of catecholamine and cortisol, which cause the rapid metabolism of high-energy storage compounds (Barton & Iwama, 1991). These catabolic processes have harmful biochemical effects on fish health and lead to slow growth (Specker & Schreck, 1980). Fish exhibit primary, secondary, and tertiary responses to stress (Barton & Iwama, 1991). The primary response is manifested in rapid changes in plasma catecholamine and corticosteroid. When these responses to stressful conditions exceed normal levels, harmful secondary and tertiary responses occur. Therefore, stress induces changes in the energetic metabolic processes, reduces growth rate, disturbs reproduction, and leads to rapid changes in flesh quality after death (Barton & Iwama, 1991). Blood chemistry serves as an indicator of an animal's physiological state and many factors (age, sex, nutrition, season, and circadian rhythm) may affect the blood chemistry. Information is available from the plasma glucose changes during starvation in the Atlantic cod, *Gadus morhua* L., the European eel, *Anguilla anguilla* (Larsson & Lewander, 1973), the pike, *Esox lucius* (Ince & Thorpe, 1976), the toadfish, *Opsanus tau* (Tashima & Cahill, 1968), the goldfish, *Carassius auratus* (Chavin & Young, 1970), and the American eel, *Anguilla rostrata* (Moon, 1983). Studies of plasma free fatty acids during starvation stress have also been performed in the European eel (Larsson & Lewander, 1973), the American eel (Moon, 1983), the rainbow trout, *Oncorhynchus mykiss* (Sumpter et al., 1991), the pike (Ince & Thorpe, 1976), and the toadfish (Tashima & Cahill, 1968). Rios et al. (2002) noted that erythrocyte senescence and hematological changes are induced by starvation in *Hoplias malabaricus*.

The Far Eastern catfish, *Silurus asotus* (Linnaeus) (order Siluriformes, family Siluridae), spreading widely throughout the Northeast Asia, is an important species in Korean freshwater aquaculture (Kim et al., 2001; Gil et al., 2017).

In this study, we tried to provide a more detailed contents, and investigated the impact of sodium nitrite on physiological response of Far Eastern catfish.

MATERIALS AND METHODS

The Far Eastern catfish, *Silurus asotus* used in this experiment was hatched in April 2015 and reared at Fishery Genetics and Breeding Sciences Laboratory of the Korea Maritime and Ocean University in Busan, Korea. Three experimental groups were established: initial, fed, and starved. All fish were fed twice daily with commercial feed (Table 1) in total 1%–

Table 1. Composition of the experimental diets used in this starvation experiment¹⁾

Nutrition	Content (%)
Crude protein	40.0
Crude fat	4.0
Crude fiber	5.0
Ash	15.0
Calcium	1.0
Phosphorus	1.0
Mineral premix ²⁾	1.0
Vitamin premix ³⁾	1.0

¹⁾ Ehwa Feed Coporation (Busan, Korea).

²⁾ Vitamin premix contained the following amount which were diluted in cellulose (g kg⁻¹ mix): L-ascorbic acid, 121.2; DL- α -tocopheryl acetate, 18.8; thiamin hydrochloride, 2.7; riboflavin, 9.1; pyridoxine hydrochloride, 1.8; niacin, 36.4; Ca-D-pantothenate, 12.7; myo-inositol, 181.8; D-biotin, 0.27; folic acid, 0.68; p-aminobenzoic acid, 18.2; menadione, 1.8; retinyl acetate, 0.73; cholecalciferol, 0.003; cyanocobalamin, 0.003.

³⁾ Mineral premix contained the following ingredients (g kg⁻¹ premix): NaCl, 43.3; MgSO₄·7H₂O, 136.5; NaH₂PO₄·2H₂O, 86.9; KH₂PO₄, 239.0; CaH₄(PO₄)·2H₂O, 135.3; ferric citrate, 29.6; ZnSO₄·7H₂O, 21.9; Ca-lactate, 304.0; CuCl, 0.2; AlCl₃·6H₂O, 0.15; KI, 0.15; Na₂Se₂O₃, 0.01; MnSO₄·H₂O, 2.0; CoCl₂·6H₂O, 1.0.

3% food of their average body weight (BW) for 2 weeks prior to initiation of experiment in March 2016. Before the starvation experiment, fish were weighed to the nearest 0.01 g using an electronic balance (JW-1, Acom, Pochen, Korea), and their body length (BL) was measured to the nearest 0.01 cm using digital Vernier calipers (CD-20CP, Mitytoyo, Kawasaki, Japan). The fed group had an average body weight (BW) of 354.1±7.74 g and average BL of 18.2±2.79 cm ($n=60$). The starved group had an average BW of 356.1±9.71 g and average BL of 18.1±2.31 cm ($n=60$).

Fish were reared in a recirculating system. 20 fish were placed each in 1.1-ton fiberglass-reinforced plastic circular tank (118 cm diameter, 100 cm depth); each experimental group consisted of three tanks of fish. Light was provided with four 40-W (5,400 K) fluorescent bulbs controlled by an electric timer, which kept the photoperiod at a 12L:12D cycle. Water temperature was controlled automatically and held at 25±0.5°C during the experimental period. During the experiment, the fed group was hand-fed two times daily (first feeding occurred between 10:00 and 11:00 and the second between 18:00 and 19:00). The starved group was starved until the end of the starvation experiment. The fed group was given feed continuously. The fish in all experiments were kept until the starved group lost vitality completely. The survival rate during the starvation experiment was calculated retroactively, with the dead fish counted every day. The aggregate survival rates of the fed group and starved group during the experimental period were measured for each of the two triplicate groups. The acute and sublethal sodium nitrite (NaNO₂) toxicity tests were conducted over period of 96 hrs. Immediately after conclusion of the starvation experiment, for the acute and NaNO₂ toxicity tests, twenty individuals (BW: 83.3±7.26 g; BL: 16.5±1.03 cm) were placed in a 150-L fiberglass-reinforced plastic tank containing 150 L of freshwater (Table 2). Nitrite (Sigma, St. Louis, MO, USA) was added to the tank to create nitrite concentration of 40–200 mg/L at intervals of 40 mg/L; nitrite-free treatment served as a control.

Table 2. Acute toxicity test condition of sodium nitrite for Far Eastern catfish, *Silurus asotus*

Test parameter ¹⁾	Conditions
Temperature (°C)	26±0.5
pH	7.3±0.45
DO (dissolved oxygen, mg/L; Saturated concentration in 26°C)	8.2
Salinity (ppt)	2.8±0.33
Ammonia (ppm)	0.02
Nitric acid (ppm)	2.4±0.12
Nitrous acid (ppm)	0.01
Conductivity (us/cm)	238

¹⁾ Temperature, pH, dissolved oxygen, and salinity were measured using an oxygen measurement electrode and a multi-data logger system (Oxyguard, Denmark). Ammonia, nitric acid, nitrous acid, and conductivity were measured using spectrophotometer (DR 2800, HACH, Loveland, Colorado, USA). The values are means of triplicate groups.

Three replications were used for each test concentration and control. Determining lethal concentration of fifty (LC₅₀) values and other related statistical analysis was based upon the method described by Peltier (1978).

Immediately after conclusion of the starvation experiment, the oxygen consumption rate was measured according to the method of Jo & Kim (1999). For the measurement of oxygen consumption rate, the respirometer chamber was utilized by a simple circulating system. The water was circulated from the reservoir (170 L) to the head tank by a circulating pump, passing by a respirometer chamber and then flowing back into the reservoir. The head tank was equipped with a temperature controller and 10 µm and 3 µm cartridge filters equipped for the filtering of particles. The flow through UV lamp was utilized for the reduction of oxygen consumption by microbes. Water from the respirometer chamber flowed into an oxygen measurement chamber. During the period of experiment, the average

water flow was 59.6 ± 0.5 L/h. The respirometer chamber was comprised of an acrylic resin box with a thickness of 8 mm; the overall dimensions of the box were 10 cm (width) \times 25 cm (length) \times 10 cm (height). A rubber pad was used as a cover for the respirometer chamber to prevent the inflow of air; a hole was made in the cover and a small valve was attached for the removal of air in the respirometer chamber. Inflow water in the respirometer chamber was diffused through a 10 mm pipe with the cap at the end, where a few holes were made. Water flowing from the respirometer chamber was flowed into the dissolved oxygen measurement chamber, the dimensions of which were 10 cm (width) \times 10 cm (length) \times 6 cm (height). The respirometer chamber consisted of three chambers, and each chamber was connected to a dissolved oxygen measurement chamber.

Dissolved oxygen measurement chambers were equipped with an oxygen probe and air in this chamber was removed by the same method as the respirometer chamber. Dissolved oxygen was measured using an oxygen measurement electrode and a multi-data logger system (Oxyguard, Denmark). Inflow and outflow dissolved oxygen of the respirometer chamber was measured by using μ Log VL 100 software at 5-minute interval over 24 hrs. Measurements of oxygen and oxygen consumption rates at each temperature were saved by the multi-data logger. After experiment started, measurement times were set at 6-hour interval over 48 hrs. Before measuring the dissolved oxygen, pH, ammonium (NH_4^+) concentration, carbon dioxide (CO_2) concentration, and respiratory frequency (gill cover movement) were measured using a counter and a digital timer. NH_4^+ and CO_2 concentrations were measured using spectrophotometer (DR 2800, HACH, Loveland, Colorado, USA). Dissolved oxygen and pH were measured using an oxygen measurement electrode and a multi-data logger system (Oxyguard, Denmark). Measurements of oxygen and oxygen consumption rates at each experimental group were saved by multi-data logger, as described by Cech (1990).

For the blood analyses of cortisol, glucose, and lactic acid, blood samples were extracted from five randomly selected fish at 0 (pre), 1, 6, 12, 24, and 48 hrs this post starvation stress test. Using syringes lined with the anticoagulant heparin blood was extracted and assayed at fixed intervals of 0, 1, 6, 12, 24, and 48 hrs from five experimental samples. Selected blood was filled into capillary tubes and analyzed after centrifuging at 200 g for 10 min. Plasma was then collected and stored in a deep freezer (SW-UF-200, Samwon Freezing Engineering, Busan, Korea) at -80°C until analysis. The cortisol concentration was measured using radioimmunoassay. Cortisol was determined in 50 μL samples using RIA kits (Coat-A-Count TKCO Cortisol RIA Kit; DPC, USA). Mixtures of sample in 100 mL antiserum were incubated for 45 min at 37°C , and then 1,000 mL separation reagent was added. The mixture was placed in a refrigerator at 4°C for 15 min and then centrifuged at $1,200 \times g$ for 15 min. Supernatant was assayed for gamma radiation using an automatic gamma counter (Cobra, Packard, USA). Plasma glucose concentration was analyzed according to methodology of Raabo and Terkildsen (1960; Kit 510, Sigma, St Louis, MO, USA), where production of H_2O_2 by glucose oxidase in the presence of *o*-dianisidine was evaluated as an absorbance increase at 450 nm. The lactic acid concentrations were analyzed using blood automatic analysis (Boehringer Mannheim Reflotron, Germany). Data were analyzed by one and two-way ANOVA with SPSS statistical package (SPSS 9.0, SPSS Inc., USA). The experiment was performed three times, and means were separated by using Duncan's multiple range test and they were considered significantly different at $p < 0.05$ (Duncan, 1955).

RESULTS

After 210 days, the starved group of Far Eastern catfish, *Silurus asotus* rapidly lost vitality, and as a result, the experiment was terminated. Fig. 1 shows changes in survival




Fig. 1. Survival rates of fed and starved Far Eastern catfish, *Silurus asotus* for 210 days experimental period. The values are means of triplicate experiments.

rate during the experimental period. Accumulated survival was 92.2±0.47% in the fed group and 74.4±2.59% in the starved group in each triplicated tanks ($p < 0.05$). Starvation resulted in retardation of growth, which has provided a good example of the reduction of the final group continuing to grow and remaining healthy (Fig. 2).

In Tables 3 and 4, and Fig. 3, the two-way ANOVA test




Fig. 2. Typical external morphology of the Far Eastern catfish, *Silurus asotus* conclusion of this starvation experiment. Initial group (a), fed group (b), and starved group (c) for 210 days period. Note the lateral ventral region in starved group is thin (Arrows in Fig. 2-c).

and measurement results of oxygen consumption rate of fed and starved groups are presented. Dissolved oxygen concentration of starved groups decreased until 48 hrs, and fed groups until 16 hrs. Table 3 shows the comparison of

Table 3. Respiratory frequency (gill cover movement) and pH values between fed group and starved group of Far Eastern catfish, *Silurus asotus*

Time (hrs)	Respiratory frequency (min ⁻¹) ¹⁾		pH ¹⁾	
	Fed group	Starved group	Fed group	Starved group
Pre-experiment	101 ^b ± 8.7	124 ^a ± 9.2	7.6 ^b ± 0.51	7.6 ^a ± 0.68
6	102 ^b ± 8.9	128 ^a ± 8.8	6.8 ^b ± 0.45	7.1 ^a ± 0.61
12	110 ^b ± 7.9	135 ^a ± 9.9	4.1 ^b ± 0.38	5.3 ^a ± 0.49
18	100 ^b ± 8.5	124 ^a ± 10.1	3.7 ^b ± 0.25	4.8 ^a ± 0.39
24	72 ^b ± 5.4	91 ^a ± 5.7	3.5 ^b ± 0.21	4.6 ^a ± 0.38
30	71 ^b ± 6.2	91 ^a ± 6.0	3.2 ^b ± 0.20	4.3 ^a ± 0.29
36	67 ^b ± 4.5	88 ^a ± 4.8	3.0 ^b ± 0.21	4.1 ^a ± 0.32
42	66 ^b ± 4.3	80 ^a ± 5.3	2.9 ^b ± 0.18	4.0 ^a ± 0.28
48	65 ^b ± 4.1	77 ^a ± 5.0	2.8 ^b ± 0.12	3.9 ^a ± 0.19

¹⁾ Respiratory frequency was measured while 1 min. Difference between fed group and starved group is significant at this level ($p < 0.05$). The values are means of triplicate experiments.

Table 4. Carbon dioxide (CO₂) and ammonium (NH₄⁺) concentration between fed group and starved group of Far Eastern catfish, *Silurus asotus*

Time (hrs)	CO ₂ concentration (mg/L) ¹⁾		Ammonium (NH ₄ ⁺) ¹⁾	
	Fed group	Starved group	Fed group	Starved group
Pre-experiment	5.1 ^b ± 0.38	5.1 ^a ± 0.41	0	0
6	14.2 ^b ± 1.28	13.0 ^a ± 1.19	0.13 ^b ± 0.012	0.08 ^a ± 0.061
12	17.2 ^b ± 1.53	14.6 ^a ± 1.21	0.17 ^b ± 0.015	0.11 ^a ± 0.018
18	18.4 ^b ± 1.67	16.4 ^a ± 1.59	0.19 ^b ± 0.017	0.14 ^a ± 0.013
24	20.0 ^b ± 1.99	18.6 ^a ± 1.87	0.23 ^b ± 0.020	0.15 ^a ± 0.014
30	20.8 ^b ± 4.89	19.4 ^a ± 1.78	0.24 ^b ± 0.019	0.16 ^a ± 0.015
36	21.2 ^b ± 2.03	20.0 ^a ± 1.90	0.26 ^b ± 0.018	0.17 ^a ± 0.016
42	21.6 ^b ± 2.10	20.4 ^a ± 1.98	0.28 ^b ± 0.209	0.18 ^a ± 0.017
48	22.0 ^b ± 2.15	20.6 ^a ± 2.12	0.29 ^b ± 0.213	0.19 ^a ± 0.017

¹⁾ Difference between fed group and starved group is significant at this level ($p < 0.05$). The values are means of triplicate experiments.

pH values and respiratory frequency (gill cover movement) between fed and starved groups. For both groups, respiratory frequency also gradually decreased until 48 hrs and pH also decreased drastically until 12 hrs, and then decreased




Fig. 3. Dissolved oxygen concentration between fed group and starved group on Far Eastern catfish, *Silurus asotus*. The values are means of triplicate experiments.

gradually until 48 hrs. The respiratory frequencies of starved group were significantly higher than those of fed group, and they decreased over the elapsed time ($p < 0.05$). Therefore, it was concluded that oxygen uptake per unit of respiratory movement was lower in starved group than in fed group. The CO₂ and NH₄⁺ concentrations increased drastically until 6 hrs, and increased gradually until 48 hrs as shown in Table 4. The CO₂ and NH₄⁺ of fed group were significantly higher than those of starved group, and they increased over the elapsed time ($p < 0.05$).

Dose-response mortality occurred in fish exposed to 120 mg/L of sodium nitrite at the fed group and 40 mg/L at the starved group (Fig. 4). Almost all fish survived 96 hrs of exposure to 0 and 40 mg/L of nitrite, whereas 75% and 25% of fish each survived 96 hrs of exposure to 150 and 200 mg/L of nitrite in fed group, and 80 and 120 mg/L of nitrite in starved group respectively. The LC₅₀, 95% confidence ranges, and experimental conditions for different exposure groups are summarized in Table 5 and Fig. 5. The LC₅₀ values for 96 hrs were 181 mg/L at the fed group and




Fig. 4. Dosage-mortality response of Far Eastern catfish, *Silurus asotus* exposed to sodium nitrite for 96 hrs. (a): fed group ($y=0.61x-61.98$, $R^2=0.98$, where y is mortality and x is sodium nitrite concentration), and (b): starved group ($y=0.45x-0.26$, $R^2=0.98$, where y is mortality and x is sodium nitrite concentration). Error bars represent the standard error of triplicate experiments ($p<0.05$).

108 mg/L at the starved group respectively.

Figs. 6, 7, and 8 are variations of hematological result for 48 hrs of exposure to sublethal sodium nitrite. In the fed group, plasma cortisol increased significantly ($p<0.05$) from 1.9 ± 0.3 $\mu\text{g/dL}$ to 30.7 ± 2.5 $\mu\text{g/dL}$ at 12 hrs, but 5.6 ± 2.9 $\mu\text{g/dL}$ at 24 hrs and 4.8 ± 0.7 $\mu\text{g/dL}$ at 48 hrs respectively. The changes for 24 hrs and 48 hrs were not




Fig. 5. Effect of starvation on LC₅₀ changes of Far Eastern catfish, *Silurus asotus* exposed to sodium nitrite at each exposure time. Fed group: $y=-2.4037x+405.74$, $R^2=0.9907$, where y is LC₅₀ and x is exposure time; starved group: $y=-2.2675x+321.69$, $R^2=0.9793$, where y is LC₅₀ and x is exposure time. Error bars represent the standard error of triplicate experiments ($p<0.05$).

significant ($p>0.05$, Fig. 6). In the starved group, plasma cortisol increased significantly ($p<0.05$) from 1.8 ± 0.3 $\mu\text{g/dL}$ to 16.7 ± 2.5 $\mu\text{g/dL}$ at 6 hrs, but 5.8 ± 2.9 $\mu\text{g/dL}$ at 12, 5.6 ± 2.9 $\mu\text{g/dL}$ 24 hrs, and 4.8 ± 0.7 $\mu\text{g/dL}$ 48 hrs respectively however, the changes for 12, 24, and 48 hrs were not significant ($p>0.05$, Fig. 6). The glucose concentrations of the two groups increased significantly from 28.5 ± 0.7 mg/dL to 242 ± 18.4 (fed) and 127 ± 3.9 mg/dL (starved) at 12 hrs ($p<0.05$ for both, Fig. 7). The glucose levels of the

Table 5. The 96 hrs LC₅₀ values of Far Eastern catfish, *Silurus asotus* exposed to sodium nitrite

Experimental group ¹⁾	96 hrs LC ₅₀ (ppm) [95% confidence ranges]	Experimental conditions ²⁾	
		DO	pH
Fed group	181.53 [171.30–199.86]	8.2±0.64	7.3±0.61
Starved group	108.92 [81.00–123.44]	8.1±0.41	7.4±0.63

¹⁾ Fed group (345 ± 20.1 mm: mean body length±SD; 562 ± 52.8 g: mean body weight±SD), starved group (243 ± 11.6 mm: mean body length±SD; 246 ± 45.8 g mean body weight±SD). The values are means of triplicate experiments.

²⁾ DO, dissolved oxygen, mg/L; Saturated concentration in 26°C.




Fig. 6. Variations in the plasma cortisol concentrations in the blood plasma of the Far Eastern catfish, *Silurus asotus*, during 48 hrs experiment. Values are means \pm SE ($n=30$). Actually $n=5$ for each experiment because the means and SE were calculated separately for each group.

fed and starved groups differed significantly from 1 hr to 48 hrs ($p < 0.05$). At 12 hrs, the cortisol and glucose levels of the fed and starved groups reached the most significant difference ($p < 0.05$). Lactic acid of fed and starved groups increased until 48 hrs, but no significant difference between fed group and starved group was found ($p > 0.05$, Fig. 8).




Fig. 7. Variations in the plasma glucose concentrations in the blood plasma of the Far Eastern catfish, *Silurus asotus*, during 48 hrs experiment. Values are means \pm SE ($n=30$). Actually $n=5$ for each experiment because the means and SE were calculated separately for each group.




Fig. 8. Variations in the lactic acid concentrations in the blood plasma of the Far Eastern catfish, *Silurus asotus*, during 48 hrs experiment. Values are means \pm SE ($n=30$). Actually $n=5$ for each experiment because the means and SE were calculated separately for each group.

DISCUSSION

Larsson and Lewander (1973) noted that many fish undergo natural periods of starvation at a particular time every year and have consequently evolved the ability to withstand the prolonged food shortages. Such periods may amount to weeks, months, or even years, and may cause extensive loss of energy stores in the body as fish consumes its own tissues to remain alive (Weatherley & Gill, 1987; Park et al., 2015). This study shows that, Far Eastern catfish, *Silurus asotus* can tolerate the long term of food shortage for 210 days from the experimental proof.

When a fish is exposed to chronic stress, its metabolic reactions are altered by changes in the hypothalamic-pituitary-adrenocortical axis (HPA axis), including the hypothalamic secretion of corticotropin-releasing hormone (CRH) through the limbic system. CRH stimulates the release of adrenocorticotropic hormone (ACTH) from the pituitary gland, which then stimulates the release of cortisol from the target organ, and the interrenal gland which promotes a metabolic stress reaction. Thus, cortisol is an important index of the stress reaction (Specker & Schreck,

1980). Severe starvation results in coma and catabolic disease of the gastrointestinal system. Basic metabolic reactions maintain energy level and body tissue (Jung et al., 2003). Guyton (1991) reported three stages of physiological change during starvation: in the first stage, the glycogen stored for urgent use is hydrolyzed, releasing glucose; in the second stage, acetyl-CoA is oversupplied, leading to acidosis; in the third stage, fish are compromised by protein exhaustion. In the process of starvation, sugar, lipid, protein, and other essential nutritional elements decrease rapidly, and finally, the whole abnormalities in immune, circulatory, and endocrine systems lead to death. In our study, dissolved oxygen concentration of starved groups decreased until 48 hrs, and fed groups until 16 hrs. Respiratory frequency also gradually decreased until 48 hrs. That is, respiratory function of all experimental groups decreased during the experimental period. Comparing the dissolved oxygen, pH, CO₂, and NH₄⁺ concentrations showed different trend, but rates of pH, CO₂, and NH₄⁺ concentrations change were similar to oxygen consumption rate. That is, the metabolic rate of all experimental groups decreased during experimental period.

The toxicity and effect of nitrite vary among fish species and depend on test conditions, such as fish size, water ion composition, and temperature (Doblender & Lackner, 1997). Chloride and other anions in water provide protective action against nitrite in active branchial uptake (Williams & Eddy, 1986); thus, small amounts of Cl⁻, for example, 1 mM, are likely to afford protection against high nitrite inputs (Eddy et al., 1983). The effect of sodium nitrite (NaNO₂) on fish is more intense in Cl⁻-poor freshwater. Thus, NaNO₂ is more toxic in freshwater than in seawater (Grosell & Jensen, 1999).

Plasma NaNO₂ accumulation probably causes methemoglobinemia and malfunction of hemopoietic activity, which are detectable effects of NaNO₂ intoxication (Costa et al., 2004). Although methemoglobinemia is not directly related to high mortality in fish exposed to NaNO₂, the

passing through of NaNO₂ into the blood stream may cause an increase in blood cell lysis, changes in the plasma electrolyte balance, and efflux of K⁺ from red blood cells, which is evident from the increase in the number of shrunken RBCs (Knudsen & Jensen, 1997; Huertas et al., 2002; Martinez & Souza, 2002; Costa et al., 2004). The dysfunctional erythrocytes may be removed from the blood circulation because of an oxygen shortage, causing reduction in the total erythrocyte counts (Park et al., 2007). For this reason, a significant difference was caused between fed and starved group in this experiment.

Plasma cortisol and plasma glucose are recognized as useful indicators of stress in fish (Park et al., 2008). Plasma cortisol and glucose levels in red drum, *Sciaenops ocellatus* simultaneously exposed to stressor were reported to be elevated (Massee et al., 1995). Barton and Iwama (1991) stated that "In general, the phenomenon that plasma cortisol concentration of fish rises by stress is first order reaction, and the phenomenon that plasma glucose concentration rises is the second-order reaction by hormone rise caused by stress". This finding has been reported in gray mullet, *Mugil cephalus* and kelp grouper, *Epinephelus bruneus* (Park et al., 2008). In this study, we have investigated the effect of starvation on Far Eastern catfish, exposed to the stress of NaNO₂. In this study, it may safely be said that we have found out a useful index for recognizing the nutritional state of Far Eastern catfish in both fed and starved group experiencing the exposure to NaNO₂.

ACKNOWLEDGEMENTS

We are grateful to anonymous reviewers for their valuable constructive comments and suggestions in improving this manuscript. We declare that all experiments in this study comply with the current laws of Korea (Ordinance of Agriculture, Food and Fisheries, No. 1 and law Regarding Experimental Animals, No. 9932).

REFERENCES

- Barton BA, Iwama GK (1991) Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Ann Rev Fish Dis* 1:3-26.
- Cech JJ (1990) Respirometry. In: Shreck CB, Moyle PB (ed), *Methods for Fish Biology*. American Fishery Society, Bethesda, pp 125-130.
- Chavin W, Young JE (1970) Factors in the determination of normal serum glucose levels of goldfish, *Carassius auratus* L. *Comp Biochem Physiol* 33:629-653.
- Costa OTF, Ferreira DJS, Mendonca FLP, Fernandes MN (2004) Susceptibility of the Amazonian fish, *Colossoma macropomum* (Serrasalminae), to short-term exposure to nitrite. *Aquaculture* 232:627-636.
- Doblender C, Lackner R (1997) Oxidation of nitrite to nitrate in isolated erythrocytes: A possible mechanism for adaptation to environmental nitrite. *Can J Fish Aquat Sci* 54:157-161.
- Duncan DB (1995) Multiple range and multiple F tests. *Biometrics* 11:1-42.
- Eddy FB, Kunzlik PA, Bath RN (1983) Uptake and loss of nitrite from the blood of rainbow trout, *Salmo gairdneri* Richardson, and Atlantic salmon, *Salmo salar* L. in fresh water and in dilute sea water. *J Fish Biol* 23:105-116.
- Gil HW, Lee TH, Han HJ, Park IS (2017) Comparative analysis of tissue and cell cycle on the far eastern catfish, *Silurus asotus* between diploid and triploid. *Dev Reprod* 21:193-204.
- Grosell M, Jensen FB (1999) NO_2^- uptake and HCO_3^- excretion in the intestine of the European flounder (*Platichthys flesus*). *J Exp Biol* 202:2103-2110.
- Guyton AC (1991) Textbook of medical physiology. In: Wonsiewicz MJ and Hallowell R (eds), *Textbook of Medical Physiology*. 8th ed. WB Saunders Company, Philadelphia p 782.
- Huertas M, Gisbert E, Rodriguez A, Cardona L, Williot P, Castello-Orvay F (2002) Acute exposure of Siberian sturgeon (*Acipenser baeri*, Brandt) yearlings to nitrite: median-lethal concentration (LC_{50}) determination, haematological changes and nitrite accumulation in selected tissues. *Aquat Toxicol* 57:257-266.
- Ince BW, Thorpe A (1976) The effects of starvation and force-feeding on the metabolism of the Northern pike, *Esox lucius* L. *J Fish Biol* 8:79-88.
- Jo J-Y, Kim YH (1999) Oxygen consumption of Far Eastern catfish, *Silurus asotus* on the different water temperatures and photoperiods. *J Korean Fish Soc* 32:56-61.
- Jung MH, Youn JM, Lee TH (2003) Effect of liriopis tuber on the change of corticosterone in mice induced by starvation stress. *Korea J Herbol* 18:279-287.
- Kim BS, Park I-S, KIM HS (2012) Effect of starvation on growth and hepatocyte nuclear size of larval haddock, *Melanogrammus aeglefinus*. *Dev Reprod* 16:107-112.
- Kim DS, Cho HJ, Park IS, Choi GC, Nam YK (2001) Cytogenetic traits and gonad development of induced triploidy in far eastern catfish (*Silurus asotus*). *Korean J Genet* 23:55-62.
- Knudsen PK, Jensen FB (1997) Recovery from nitrite-induced methaemoglobinaemia and potassium balance disturbances in carp. *Fish Physiol Biochem* 16:1-10.
- Larsson A, Lewander K (1973) Metabolic effects of starvation in the eel, *Anguilla anguilla* L. *Comp Biochem Physiol A* 44:367-374.
- Martinez CBR, Souza MM (2002) Acute effects of nitrite on ion regulation in two neotropical fish species. *Comp Biochem Physiol* 133A:151-160.
- Massee KC, Rust MB, Hardy RW, Stickney RR (1995) The effectiveness of tricaine, quinaldine sulfate and metomidate as anesthetics for larval fish. *Aquaculture* 134:351-359.
- Mehner T, Wieser W (1994) Energetics and metabolic correlates of starvation in juvenile perch, *Perca fluviatilis*.

- J Fish Biol 45:325-333.
- Moon TW. (1983) Metabolic reserves and enzyme activities with food deprivation in immature American eels, *Anguilla rostrata* (Lesueur). Can J Zool 61:802-811.
- Mustafa S, Mittal A (1982) Protein, RNA and DNA levels in liver and brain of starved catfish, *Clarias batrachus*. Jpn J Ichthyol 28:396-400.
- Park IS, Choi HJ, Noh CH, Myoung JG, Park HJ, Goo IB (2013) Induced morphological changes in larval rock bream, *Oplegnathus fasciatus*, under starvation. Dev Reprod 17:399-407.
- Park IS, Gil HW, Yoo GY, Oh JS (2015) Effects of starvation in rock bream, *Oplegnathus fasciatus* and olive flounder, *Paralichthys olivaceus*. Dev Reprod 19:97-109.
- Park IS, Im JH, Ryu DK, Nam YK, Kim DS (2001) Effect of starvation on morphometric changes in *Rhynchocypris oxycephalus* (Sauvage and Dabry). J Appl Ichthyol 17:277-281.
- Park IS, Lee JH, Hur JW, Song YC, Na HC, Noh CH (2007) Acute toxicity and sublethal effects of nitrite on selected hematological parameters and tissues in dark-banded rockfish, *Sebastes inermis*. J World Aquac Soc 38:188-199.
- Park MO, Hur WJ, Im SY, Seol DW, Lee JH, Park IS (2008) Anaesthetic efficacy and physiological responses to clove oil-anaesthetized kelp grouper *Epinephelus bruneus*. Aquac Res 39: 877-884.
- Peltier W (1978) Methods for measuring the acute toxicity for effluents to aquatic organisms. Environmental Protection Agency, Office of Research and Development, National Technical Information Service, Springfield, Virginia, USA.
- Rios FS, Kalinin AL, Rantin FT (2002) The effects of long-term food deprivation on respiration and haematology of the neotropical fish *Hoplias malabaricus*. J Fish Biol 61:85-95.
- Specker JL, Schreck CB (1980) Stress responses to transportation and fitness for marine survival in coho salmon, *Oncorhynchus kisutch* smolts. Can J Fish Aquat Sci 37:765-769.
- Sumpter JP, Le Bail PY, Pickering AD, Pottinger TG, Carragher JF (1991) The effect of starvation on growth and plasma growth hormone concentrations of rainbow trout, *Oncorhynchus mykiss*. Gen Comp Endocrinol 83: 94-102.
- Tashima L, Cahill GF (1968) Effects of insulin in the toadfish, *Opsanus tau*. Gen Comp Endocrinol 11:262-271.
- Weatherley AH, Gill HS (1987) The Biology of Fish Growth. 4. Protein, Lipid and Caloric Contents. London, Academic Press. pp 139-146.
- Williams EM, Eddy FB (1986) Chloride uptake in freshwater teleosts and its relationship to nitrite uptake and toxicity. J Comp Physiol B 156:867-872.