

Role of Type 1 Inositol 1,4,5-triphosphate Receptors in Mammalian Oocytes

†Sook Young Yoon

Fertility Center of CHA Gangnam Medical Center, CHA University, Seoul 06125, Korea

ABSTRACT : The ability of oocytes to undergo normal fertilization and embryo development is acquired during oocyte maturation which is transition from the germinal vesicle stage (GV), germinal vesicle breakdown (GVBD) to metaphase of meiosis II (MII). Part of this process includes redistribution of inositol 1, 4, 5-triphosphate receptor (IP3R), a predominant Ca^{2+} channel on the endoplasmic reticulum membrane. Type 1 IP3R (IP3R1) is expressed in mouse oocytes dominantly. At GV stage, IP3R1 are arranged as a network throughout the cytoplasm with minute accumulation around the nucleus. At MII stage, IP3R1 diffuses to the entire cytoplasm in a more reticular manner, and obvious clusters of IP3R1 are observed at the cortex of the egg. This structural reorganization provides acquisition of $[Ca^{2+}]_i$ oscillatory activity during fertilization. In this review, general properties of IP3R1 in somatic cells and mammalian oocyte are introduced.

Key words : IP3R1, Oocytes, Fertilization, PLCzeta

Increase of intracellular free calcium concentration ($[Ca^{2+}]_i$) is monitored in various cellular response to stimuli (Berridge et al., 2000). $[Ca^{2+}]_i$ increases is observed in most of cellular response such as secretion and contraction, proliferation, differentiation, fertilization, development and cell death (Berridge et al., 2003). Fertilization of mammalian oocytes, one of the cellular event, is controlled by increase in $[Ca^{2+}]_i$ with oscillation pattern that persists for several hours from sperm-egg fusion (Miyazaki et al., 1993). These $[Ca^{2+}]_i$ oscillations are responsible for the initiation of fertilization event including, cortical granule exocytosis to block to polyspermy, resumption of meiosis, recruitment of maternal mRNAs and activation of embryonic genome, and further embryonic development (Ajduk et al., 2008). Most Ca^{2+} -increase during fertilization origi-

nates from intracellular calcium store via 1, 4, 5 inositol triphosphate (IP3) receptors (IP3R) on the endoplasmic reticulum membrane. IP3 is a widespread signaling molecule produced as a result of hydrolysis of phosphatidylinositol 4, 5-bisphosphate (PIP₂) by phospholipase C (PLC) (Lee et al., 2006b). Egg has reported to express all three isoforms of IP3R, but type 1 IP3R (IP3R1) is present significantly larger amount in mammalian eggs (He et al., 1999). This review is to introduce the general information of IP3R and role of IP3R in mammalian egg fertilization.

$[Ca^{2+}]_i$ OSCILLATION DURING FERTILIZATION

Fertilization is a process which is regulated by IP3-

Manuscript received January 10, 2019, Received in revised form February 17, 2019, Accepted February 27, 2019

† Corresponding Author : Sook Young Yoon, Fertility Center of CHA Gangnam Medical Center, CHA University, Seoul 06125, Korea. Tel: +82-2-3468-2841, E-mail: syyoon11@cha.ac.kr



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

mediated Ca^{2+} -release from intracellular Ca^{2+} -store. In mammalian oocytes, the sperm factor (SF, now it has been as known phospholipase C zeta, $\text{PLC}\zeta$) triggers activation of phosphoinositide (PI) pathway that produces IP3 and 1, 2-diacylglycerol (DAG) by hydrolysis of PIP2. Increase intracellular IP3 concentration is in charge for mediating Ca^{2+} -release from endoplasmic reticulum (ER) via IP3 receptor (Fig. 1) (Miyazaki et al., 1993; Malcuit et al., 2005; Malcuit et al., 2006; Yoon, 2011). Microinjection of 18A10 antibody, which recognizes an epitope to the Ca^{2+} channel region in the C-terminus of IP3R1, or IP3R antagonist inhibits $[\text{Ca}^{2+}]_i$ oscillation in fertilized oocytes (Miyazaki et al., 1992). Also, adenophostin A, agonist of IP3R, or thimerosal, a thiol oxidizing agent, induced $[\text{Ca}^{2+}]_i$ oscillation and further egg activation process in mature oocytes without sperm (Fissore et al., 1992; Jellerette et al., 2000).

GENERAL STRUCTURE OF IP3R1

IP3, as an intracellular second messenger, mediating

Ca^{2+} -signalling in cellular response was first reported in neutrophils (Prentki et al., 1984). Autoradiographic studies with $[\text{H}^3]$ IP3 represented high densities of IP3 binding in cerebellum with 100–300 times higher than in peripheral tissues (Worley et al., 1987). The purification of IP3R was reported 260 KDa protein band by electrophoretic analysis, purified protein presented selectivity for IP3 over other inositol phosphates (Supattapone et al., 1988). Three isoforms of IP3R (type 1, type 2, and type 3) have been identified by cDNA cloning. Most cells express at least one isoform and type 1 IP3R (IP3R1) is the most widely expressed in all cell types and all developmental stages (Marks, 1997).

The three IP3R isoforms demonstrate different IP3-binding affinities and sensitivity. The overall domain structure of IP3R contains a cytoplasmic N-terminus, IP3 binding domain, regulatory or modulatory domain which has multiple regulatory sites for Ca^{2+} , calmodulin dependent protein kinase, and other regulatory protein (Fig. 2). The C-terminus is channel domain including six transmembrane




Fig. 1. Role of IP3R1 during fertilization in mammalian oocytes. ① $\text{PLC}\zeta$ hydrolyzes PIP2 to DAG (③) and IP3, ② IP3 binds to IP3R1 on the endoplasmic reticulum, ④ Activated IP3R1 releases Ca^{2+} from ER to cytoplasm, ⑤ Ca^{2+} oscillation in the cytoplasm.



Fig. 2. Overall structure of IP3R.

domains (TMD), which form Ca^{2+} -release channels on the endoplasmic reticulum membrane (Lee et al., 2006b).

Using electron microscopy analysis, IP3R was observed tetrameric structure with reversible transition between two distinct structure, windmill and square structure (Hamada et al., 2002).

ACTIVATION OF IP3R

Ca^{2+} -release through IP3R is known to be modulated by posttranslational modification of IP3R including, binding with IP3 and Ca^{2+} , clustering of IP3R, phosphorylation / dephosphorylation, and spatial distribution in the cytoplasm.

Activation of IP3R is begun by IP3 binding to the IP3 binding domain (amino acid 226–576) and Ca^{2+} ion permeating. At least four IP3 binding sites of the tetrameric structure of IP3R are essential to activate for IP3R (Taylor & Tovey, 2010). In addition, cytosolic Ca^{2+} was shown to regulate IP3, however, the effects of Ca^{2+} were biphasic. The modest increase of intracellular Ca^{2+} induces the enhancing response to IP3, while higher increase have inhibitory effects in IP3R1 through patch clamp electrophysiology of isolated *Xenopus* oocyte nuclei membrane (Mak et al., 2003). At least eight Ca^{2+} -binding sites have been identified in IP3R1, Ca^{2+} promotes the transition from the square (or mushroom-like structure) to the windmill structure with relocation of four peripheral IP3 binding domains (Hamada et al., 2003).

1. Clustering of IP3R

A common consequence of activation of IP3R is the formation of IP3R clusters. IP3R clustering in somatic cells has been known to temporarily concur with increases in the cytosolic IP3 rather than with $[\text{Ca}^{2+}]_i$ (Lee et al.,

2006b). Green fluorescence protein-tagged IP3R1 was expressed on the rat hippocampal neuron *in vitro*, and its lateral diffusion was regulated by actin filaments and 4.1N proteins, cytoskeletal-associated proteins enriched in neuron (Fukatsu et al., 2004). Single IP3R response first, then clustered IP3Rs open together with local Ca^{2+} -puff, then all puffs lead to a global regenerative Ca^{2+} waves that spread out through whole cytoplasm in general somatic cells (Rahman, 2009, 2012). This IP3R clustering induced by IP3 production in the absence of a significant Ca^{2+} increase, but elevated Ca^{2+} without IP3 production did not induce IP3R clustering (Tateishi et al., 2005).

2. Phosphorylation/dephosphorylation of IP3R in general cellular response

IP3R isoforms have multiple phosphorylation consensus sites and several docking sites for protein kinase and phosphatases on their sequences. Today at least 12 different protein kinases are investigated.

1) Protein kinase A (PKA)

PKA phosphorylates two distinct sites on IP3R1 (S¹⁵⁸⁸ and S¹⁷⁵⁵) in the regulatory region, and phosphorylation of this sites regulates channel open probability of IP3R1 in whole cell patch clamp study (Wagner et al., 2008). Association of IP3R1 with protein phosphatase 1 alpha facilitates dephosphorylation of PKA mediated phosphorylation (Tang et al., 2003). Also, involvement of PKA to IP3R1 activation is mediated by leucine / isoleucine zipper domain in the regulatory region (Tu et al., 2004).

2) Protein kinase G (PKG)

Another protein kinase G is shown restricted tissue distribution than PKA, could phosphorylate IP3R1 (Ser⁶⁹⁶) with

IP3R-associated cGMP kinase substrate (IRAG) in microsomal smooth muscle membranes (Schlossmann et al., 2000).

3) Ca²⁺/CaM-dependent protein kinase II

Ca²⁺/CaM-dependent protein kinase II (CaM KII) is a serine / threonine protein kinase expressed as an 8–12 monomers in most tissues, and highly sensitive to Ca²⁺ and calmodulin. Also it has been shown as Ca²⁺-oscillation decoder in various cellular responses (Dupont & Goldbeter, 1998). All three different CaM KII inhibitors (KN-93, KN-62, and Autocamtide - 2 - Related Inhibitory Peptide (AIP)) inhibited ATP-mediated [Ca²⁺]_i transient in calf pulmonary artery endothelial *in vitro* (Aromolaran & Blatter, 2005). Colocalization of CaM KII and IP3R have reported in several tissues, this interaction induced phosphorylation of IP3R within N-terminal region. Besides, this phosphorylation of IP3R significantly decreases the open probability in lipid bilayers. It may suggest CaM KII acts on Ca²⁺ oscillations negatively (Zima et al., 2007).

4) Protein kinase C (PKC)

The activity of conventional PKCs, including α , β , and γ , depend on Ca²⁺ and diacylglycerol. Also, PKC activation leads to changes of subcellular localization of IP3Rs in many cell types (Vermassen et al., 2003). Phosphorylation sites for PKC on IP3R is [R/K]-X-[S/T]-X-[R/K], and counteracts by phosphatase calcineurin (Cameron et al., 1995). In addition, there are several proteins to interact with PKC for IP3R activation, FKBP12, mTOR and RACK1.

5) Protein kinase B (PKB)

All three IP3R isoforms contain an R-X-R-X-X-[S/T] consensus site for PKB in their C-terminal. PKB has survival effects on apoptosis by inhibiting IP3-induced Ca²⁺ release on S²⁶⁸¹ (Joseph & Hajnoczky, 2007).

6) Cyclin-dependent protein kinases (CDKs)

Cyclin-dependent kinases (CDKs) are a family of protein

kinases and involved in cell cycle, including extracellular signal regulated kinase (ERK), and polo like kinase.

A well investigated cellular system, mammalian oocytes, was shown these protein kinases activity changes during oocyte maturation. Increase of [Ca²⁺]_i during fertilization has been reported IP3R mediated Ca²⁺ release with single, large Ca²⁺ increase followed by Ca²⁺ oscillation for several hours until pronuclear formation. These Ca²⁺ changes depend on IP3R activity through protein kinase activity. CDKs (*cdc2*) assemble with the regulatory protein cyclin B to form the maturation promoting factor, which is major protein for starting of oocyte maturation (Polanski et al., 2012). CDK1 phosphorylates IP3R1 at S⁴²¹, in IP3 binding domain and T⁷⁹⁹ in regulatory domain. Both sites are conserved in IP3R1 from human to *Drosophila* (Lee et al., 2006b). During oocyte maturation, IP3R1 undergoes cell cycle-dependent phosphorylation in mouse oocytes, M-phase kinases associates as regulators of IP3R1 in mammalian egg (Jellerette et al., 2004). CDK1/cyclin B mediated IP3R1 phosphorylation at T⁷⁹⁹ increases IP3 binding activity, resulting in an increase Ca²⁺-releasing activity.

ERK1 and ERK2 are cell cycle dependent kinases and phosphorylate an P-X-[S/T]-P. In mouse IP3R1 has three ERK1/2 phosphorylation sites for ERK1/2 at S⁴³⁶, T⁹⁴⁵ and S¹⁷⁶⁵. From *in vitro* phosphorylation studies represent that ERK2 phosphorylates IP3R1 at S⁴³⁶, that mutation (S436A) abrogates IP3R1 phosphorylation (Lee et al., 2006a).

7) Tyrosine kinases

Phosphorylation of tyrosine residues of IP3R1 was observed in T-cell receptor activation, the IP3R phosphorylation is induced by tyrosine kinase, Fyn at Y³⁵³. This phosphorylation was reduced in thymocytes in *fyn*^{-/-} mice (Jayaraman et al., 1996). In T-cell activation, IP3R phosphorylation by Fyn induces increase of affinity for IP3 to 5 times, which means that IP3R1 can continue to release Ca²⁺ from intracellular Ca²⁺ store during the reducing of intracellular IP3 (Cui et al., 2004).

8) Anti-apoptotic Bcl-2 protein

B-cell lymphoma 2 (Bcl-2), encoded in humans by the *Bcl2* gene, is a proapoptotic and anti-apoptotic protein with four BH domain and embedded in the endoplasmic reticulum, nuclear envelope, and mitochondrial outer membrane (Greenberg et al., 2014). Bcl-2 proteins have been associated on PKA mediated phosphorylation.

CHANGES OF IP3R1 DURING OOCYTE MATURATION IN MAMMALS

Multiple changes in Ca^{2+} -response mechanism via IP3R1 including IP3R1 clustering on the plasma membrane (Mehlmann & Kline, 1994; Mehlmann et al., 1996; Machaca, 2004; Sun et al., 2011), phosphorylation of IP3R1 (Vanderheyden et al., 2009; Wakai et al., 2012; Zhang et al., 2015), have been described during oocyte maturation.

1. Type of IP3R in mammalian oocytes

As IP3R1 is observed in whole cytoplasm, 1.9 fold maturation associated increase was indicated by western blot analysis, in contrast, the type 2 and the type 3 IP3R were detected lesser in oocytes than IP3R1 (Fissore et al., 1999).

2. Mass of IP3R1 protein in oocytes

During the mouse oocyte maturation, it has known to increase an sensitivity to IP3 mediated release of intracellular calcium. 1.5 to 2 fold maturation associated increase in the mass of IP3R1 (Fissore et al., 1999) stands for this increase in IP3 sensitivity, and administrated dsRNA corresponding to the IP3R1 sequence into germinal vesicle (GV) immature oocytes and resulted in the mass of IP3R1 protein and significantly decreased Ca^{2+} transients in these oocytes than controls (Xu et al., 2003).

3. Subcellular localization, clustering and regulation of IP3R1 function in mammalian egg

The sensitivity of Ca^{2+} release has known to increase

during oocyte maturation from GV stage to MII stage (Mehlmann & Kline, 1994; Mehlmann et al., 1996). In immature GV oocytes showed homogeneous distribution of IP3R1 on entire cytoplasm except geminal vesicle. However, during oocyte maturation, IP3R1 reorganized their localization and clustering itself. Most of the IP3R1 distributed in whole cytoplasm with cortical clusters near plasma membrane in large clusters, 1-2 μm in diameter (yellow arrow in Fig. 3) (Mehlmann et al., 1996). Inhibition of this clustering by MEK-specific inhibitor, U0126, reduced phosphorylation of IP3R1 and calcium release during fertilization (Lee et al., 2006a; Ito et al., 2008). Changes in IP3R1 sensitivity may underline the changes of the spatio-temporal Ca^{2+} responses during oocyte maturation. The conductivity of IP3R1 to Ca^{2+} in response to increase in IP3 is enhanced at mature MII oocytes (Machaca, 2007; Ullah et al., 2007). The Ca^{2+} transient by PLC ζ in GV showed very small and narrow spike, however, that in mature MII oocytes showed big and long (4–5 mins) spike




Fig. 3. IP3R1 distribution in GV, GVBD and MII mouse oocytes. GVBD oocytes were cultured for 4 hr, or 8 hr *in vitro*. Arrows indicate cortical clusters more than 1 μm in diameter. Scale bar=10 μm . GV, germinal vesicle ; GVBD, germinal vesicle breakdown.

in mouse oocytes (Machaca, 2007). Also Ca^{2+} -oscillations in response to sperm of immature oocytes were maximum three much smaller transients, which ceased after 1 hr. But the oscillation of Ca^{2+} in mature oocytes by sperm continued for 4 to 5 hrs, near pronuclear (PN) formation (Jones et al., 1995a; Jones et al., 1995b). These sensitivity of Ca^{2+} response is controlled by M phase kinase during oocyte maturation with cell cycle dependent (Jones et al., 1995a; Jellerette et al., 2004). During oocyte maturation GV immature oocytes have nonphosphorylated form of IP3R1, on the other hands, MII mature oocytes have phosphorylated IP3R1 on S⁴³⁶ by MAPK, and S⁴²¹, T⁷⁹⁹ (Lee et al., 2006b; Zhang et al., 2015).

4. IP3R1 degradation during fertilization

In mammalian oocytes, fertilization mediated Ca^{2+} -oscillations are very active at the beginning of fertilization, which is ceased as zygotes come close to the pronuclear stage. Further, more than 50% of IP3R1 mass is down-regulated by PN stage (Brind et al., 2000; Jellerette et al., 2000; Jones, 2005). These IP3R1 degradation requires continuous identical and persistently elevated levels of IP3 by PLC ζ (Jellerette et al., 2000). Also continuous degradation of the IP3R1 results in Ca^{2+} -oscillation with reduced periodicity (Lee et al., 2010). Degradation of IP3R1 is mediated by ubiquitin-proteasome pathway after IP3 binding in intact mammalian cells (Alzayady & Wojcikiewicz, 2005).

CONCLUSION

The generating of Ca^{2+} -release during fertilization provides the widespread intracellular signaling for egg activation and further embryonic development. In mammalian oocytes, Ca^{2+} oscillation during fertilization persists for several hours until the pronuclear formation. Here we interpret general characteristic of IP3R1 and the role of IP3R1 during fertilization. Further studies should be inves-

tigated the molecular mechanism that controls the activation and degradation of IP3R1 in egg activation and further embryonic development. It is hoped that this work will help the understanding of physiological problem in infertility or subfertility patients.

ACKNOWLEDGMENTS

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2014R1A1A2056914).

REFERENCES

- Ajduk A, Malagocki A, Maleszewski M (2008) Cytoplasmic maturation of mammalian oocytes: Development of a mechanism responsible for sperm-induced Ca^{2+} oscillations. *Reprod Biol* 8:3-22.
- Alzayady KJ, Wojcikiewicz RJH (2005) The role of Ca^{2+} in triggering inositol 1,4,5-trisphosphate receptor ubiquitination. *Biochem J* 392:601-606.
- Aromolaran AAS, Blatter LA (2005) Modulation of intracellular Ca^{2+} release and capacitative Ca^{2+} entry by CaMKII inhibitors in bovine vascular endothelial cells. *Am J Physiol Cell Physiol* 289:C1426-C1436.
- Berridge MJ, Bootman MD, Roderick HL (2003) Calcium signalling: Dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* 4:517-529.
- Berridge MJ, Lipp P, Bootman MD (2000) The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* 1:11-21.
- Brind S, Swann K, Carroll J (2000) Inositol 1,4,5-trisphosphate receptors are downregulated in mouse oocytes in response to sperm or adenophostin A but not to increases in intracellular Ca^{2+} or egg activation. *Dev Biol* 223:251-265.
- Cameron AM, Steiner JP, Roskams AJ, Ali SM, Ronnett

- GV, Snyder SH (1995) Calcineurin associated with the inositol 1,4,5-trisphosphate receptor-FKBP12 complex modulates Ca^{2+} flux. *Cell* 83:463-472.
- Cui J, Matkovich SJ, deSouza N, Li S, Rosemblyt N, Marks AR (2004) Regulation of the type 1 inositol 1,4,5-trisphosphate receptor by phosphorylation at tyrosine 353. *J Biol Chem* 279:16311-16316.
- Dupont G, Goldbeter A (1998) CaM kinase II as frequency decoder of Ca^{2+} oscillations. *Bioessays* 20:607-610.
- Fissore RA, Dobrinsky JR, Balise JJ, Duby RT, Robl JM (1992) Patterns of intracellular Ca^{2+} concentrations in fertilized bovine eggs. *Biol Reprod* 47:960-969.
- Fissore RA, Longo FJ, Anderson E, Parys JB, Ducibella T (1999) Differential distribution of inositol trisphosphate receptor isoforms in mouse oocytes. *Biol Reprod* 60:49-57.
- Fukatsu K, Bannai H, Zhang S, Nakamura H, Inoue T, Mikoshiba K (2004) Lateral diffusion of inositol 1,4,5-trisphosphate receptor type 1 is regulated by actin filaments and 4.1N in neuronal dendrites. *J Biol Chem* 279:48976-48982.
- Greenberg EF, Lavik AR, Distelhorst CW (2014) Bcl-2 regulation of the inositol 1,4,5-trisphosphate receptor and calcium signaling in normal and malignant lymphocytes: Potential new target for cancer treatment. *Biochim Biophys Acta* 1843:2205-2210.
- Hamada K, Miyata T, Mayanagi K, Hirota J, Mikoshiba K (2002) Two-state conformational changes in inositol 1,4,5-trisphosphate receptor regulated by calcium. *J Biol Chem* 277:21115-21118.
- Hamada K, Terauchi A, Mikoshiba K (2003) Three-dimensional rearrangements within inositol 1,4,5-trisphosphate receptor by calcium. *J Biol Chem* 278:52881-52889.
- He CL, Damiani P, Ducibella T, Takahashi M, Tanzawa K, Parys JB, Fissore RA (1999) Isoforms of the inositol 1,4,5-trisphosphate receptor are expressed in bovine oocytes and ovaries: The type-1 isoform is down-regulated by fertilization and by injection of adenosine phosphatase. *Biol Reprod* 61:935-943.
- Ito J, Yoon SY, Lee B, Vanderheyden V, Vermassen E, Wojcikiewicz R, Alfandari D, De Smedt H, Parys JB, Fissore RA (2008) Inositol 1,4,5-trisphosphate receptor 1, a widespread Ca^{2+} channel, is a novel substrate of polo-like kinase 1 in eggs. *Dev Biol* 320:402-413.
- Jayaraman T, Ondrias K, Ondria E, Marks AR (1996) Regulation of the inositol 1,4,5-trisphosphate receptor by tyrosine phosphorylation. *Science* 272:1492-1494.
- Jellerette T, He CL, Wu H, Parys JB, Fissore RA (2000) Down-regulation of the inositol 1,4,5-trisphosphate receptor in mouse eggs following fertilization or parthenogenetic activation. *Dev Biol* 223:238-250.
- Jellerette T, Kurokawa M, Lee B, Malcuit C, Yoon SY, Smyth J, Vermassen E, De Smedt H, Parys JB, Fissore RA (2004) Cell cycle-coupled $[\text{Ca}^{2+}]_i$ oscillations in mouse zygotes and function of the inositol 1,4,5-trisphosphate receptor-1. *Dev Biol* 274:94-109.
- Jones KT (2005) Mammalian egg activation: From Ca^{2+} spiking to cell cycle progression. *Reproduction* 130:813-823.
- Jones KT, Carroll J, Merriman JA, Whittingham DG, Kono T (1995a) Repetitive sperm-induced Ca^{2+} transients in mouse oocytes are cell cycle dependent. *Development* 121:3259-3266.
- Jones KT, Carroll J, Whittingham DG (1995b) Ionomycin, thapsigargin, ryanodine, and sperm induced Ca^{2+} release increase during meiotic maturation of mouse oocytes. *J Biol Chem* 270:6671-6677.
- Joseph SK, Hajnoczky G (2007) IP3 receptors in cell survival and apoptosis: Ca^{2+} release and beyond. *Apoptosis* 12:951-968.
- Lee B, Vermassen E, Yoon SY, Vanderheyden V, Ito J, Alfandari D, De Smedt H, Parys JB, Fissore RA (2006a) Phosphorylation of IP3R1 and the regulation of $[\text{Ca}^{2+}]_i$ responses at fertilization: A role for the MAP kinase pathway. *Development* 133:4355-4365.

- Lee B, Yoon SY, Fissore RA (2006b) Regulation of fertilization-initiated $[Ca^{2+}]_i$ oscillations in mammalian eggs: A multi-pronged approach. *Semin Cell Dev Biol* 17:274-284.
- Lee B, Yoon SY, Malcuit C, Parys JB, Fissore RA (2010) Inositol 1,4,5-trisphosphate receptor 1 degradation in mouse eggs and impact on $[Ca^{2+}]_i$ oscillations. *J Cell Physiol* 222:238-247.
- Machaca K (2004) Increased sensitivity and clustering of elementary Ca^{2+} release events during oocyte maturation. *Dev Biol* 275:170-182.
- Machaca K (2007) Ca^{2+} signaling differentiation during oocyte maturation. *J Cell Physiol* 213:331-340.
- Mak DOD, McBride SMJ, Petrenko NB, Foskett JK (2003) Novel regulation of calcium inhibition of the inositol 1,4,5-trisphosphate receptor calcium-release channel. *J Gen Physiol* 122:569-581.
- Malcuit C, Knott JG, He C, Wainwright T, Parys JB, Robl JM, Fissore RA (2005) Fertilization and inositol 1,4,5-trisphosphate (IP3)-induced calcium release in type-1 inositol 1,4,5-trisphosphate receptor down-regulated bovine eggs. *Biol Reprod* 73:2-13.
- Malcuit C, Kurokawa M, Fissore RA (2006) Calcium oscillations and mammalian egg activation. *J Cell Physiol* 206:565-573.
- Marks AR (1997) Intracellular calcium-release channels: Regulators of cell life and death. *Am J Physiol* 272:H597-H605.
- Mehlmann LM, Kline D (1994) Regulation of intracellular calcium in the mouse egg: Calcium release in response to sperm or inositol trisphosphate is enhanced after meiotic maturation. *Biol Reprod* 51:1088-1098.
- Mehlmann LM, Mikoshiba K, Kline D (1996) Redistribution and increase in cortical inositol 1,4,5-trisphosphate receptors after meiotic maturation of the mouse oocyte. *Dev Biol* 180:489-498.
- Miyazaki S, Shirakawa H, Nakada K, Honda Y (1993) Essential role of the inositol 1,4,5-trisphosphate receptor/ Ca^{2+} release channel in Ca^{2+} waves and Ca^{2+} oscillations at fertilization of mammalian eggs. *Dev Biol* 158:62-78.
- Miyazaki S, Yuzaki M, Nakada K, Shirakawa H, Nakanishi S, Nakade S, Mikoshiba K (1992) Block of Ca^{2+} wave and Ca^{2+} oscillation by antibody to the inositol 1,4,5-trisphosphate receptor in fertilized hamster eggs. *Science* 257:251-255.
- Polanski Z, Homer H, Kubiak JZ (2012) Cyclin B in mouse oocytes and embryos: Importance for human reproduction and aneuploidy. *Results Probl Cell Differ* 55:69-91.
- Prentki M, Wollheim CB, Lew PD (1984) Ca^{2+} homeostasis in permeabilized human neutrophils. Characterization of Ca^{2+} -sequestering pools and the action of inositol 1,4,5-triphosphate. *J Biol Chem* 259:13777-13782.
- Rahman T (2012) Dynamic clustering of IP3 receptors by IP3. *Biochem Soc Trans* 40:325-330.
- Rahman TU, Skupin A, Falcke M, Taylor CW (2009) Clustering of InsP3 receptors by InsP3 retunes their regulation by InsP3 and Ca^{2+} . *Nature* 458:655-659.
- Schlossmann J, Ammendola A, Ashman K, Zong X, Huber A, Neubauer G, Wang GX, Allescher HD, Korth M, Wilm M, Hofmann F, Ruth P (2000) Regulation of intracellular calcium by a signalling complex of IRAG, IP3 receptor and cGMP kinase I β . *Nature* 404:197-201.
- Sun L, Yu F, Ullah A, Hubrack S, Daalis A, Jung P, Machaca K (2011) Endoplasmic reticulum remodeling tunes IP3-dependent Ca^{2+} release sensitivity. *PLOS ONE* 6:e27928.
- Supattapone S, Worley PF, Baraban JM, Snyder SH (1988) Solubilization, purification, and characterization of an inositol trisphosphate receptor. *J Biol Chem* 263:1530-1534.
- Tang TS, Tu H, Wang Z, Bezprozvanny I (2003) Modulation of type 1 inositol (1,4,5)-trisphosphate receptor function by protein kinase A and protein phosphatase.

- phatase 1alpha. *J Neurosci* 23:403-415.
- Tateishi Y, Hattori M, Nakayama T, Iwai M, Bannai H, Nakamura T, Michikawa T, Inoue T, Mikoshiba K (2005) Cluster formation of inositol 1,4,5-trisphosphate receptor requires its transition to open state. *J Biol Chem* 280:6816-6822.
- Taylor CW, Tovey SC (2010) IP(3) receptors: Toward understanding their activation. *Cold Spring Harb Perspect Biol* 2:a004010.
- Tu H, Tang TS, Wang Z, Bezprozvanny I (2004) Association of type 1 inositol 1,4,5-trisphosphate receptor with AKAP9 (Yotiao) and protein kinase A. *J Biol Chem* 279:19375-19382.
- Ullah G, Jung P, Machaca K (2007) Modeling Ca^{2+} signaling differentiation during oocyte maturation. *Cell Calcium* 42:556-564.
- Vanderheyden V, Wakai T, Bultynck G, De Smedt H, Parys JB, Fissore RA (2009) Regulation of inositol 1,4,5-trisphosphate receptor type 1 function during oocyte maturation by MPM-2 phosphorylation. *Cell Calcium* 46:56-64.
- Vermassen E, Van Acker K, Annaert WG, Himpens B, Callewaert G, Missiaen L, De Smedt H, Parys JB (2003) Microtubule-dependent redistribution of the type-1 inositol 1,4,5-trisphosphate receptor in A7r5 smooth muscle cells. *J Cell Sci* 116:1269-1277.
- Wagner LE, Joseph SK, Yule DI (2008) Regulation of single inositol 1,4,5-trisphosphate receptor channel activity by protein kinase A phosphorylation. *J Physiol* 586:3577-3596.
- Wakai T, Vanderheyden V, Yoon SY, Cheon B, Zhang N, Parys JB, Fissore RA (2012) Regulation of inositol 1,4,5-trisphosphate receptor function during mouse oocyte maturation. *J Cell Physiol* 227:705-717.
- Worley PF, Baraban JM, Colvin JS, Snyder SH (1987) Inositol trisphosphate receptor localization in brain: Variable stoichiometry with protein kinase C. *Nature* 325:159-161.
- Xu Z, Williams CJ, Kopf GS, Schultz RM (2003) Maturation-associated increase in IP3 receptor type 1: Role in conferring increased IP3 sensitivity and Ca^{2+} oscillatory behavior in mouse eggs. *Dev Biol* 254:163-171.
- Yoon SY, Kang DW (2011) Production of intracellular calcium oscillation by phospholipase C zeta activation in mammalian eggs. *Dev Reprod* 15:197-204.
- Zhang N, Yoon SY, Parys JB, Fissore RA (2015) Effect of M-phase kinase phosphorylations on type 1 inositol 1,4,5-trisphosphate receptor-mediated Ca^{2+} responses in mouse eggs. *Cell Calcium* 58:476-488.
- Zima AV, Bare DJ, Mignery GA, Blatter LA (2007) IP3-dependent nuclear Ca^{2+} signalling in the mammalian heart. *J Physiol* 584:601-611.