

Expressional Evaluation of C/EBP Family, SREBP1, and Steroid Hormone Receptors in the Epididymal Fat of Postnatally Developing Mouse

Yong-Seung Lee¹ and [†]Ki-Ho Lee²

¹Dept. of Biological Science, Sungkyunkwan University, Suwon 16419, Korea

²Dept. of Biochemistry and Molecular Biology, College of Medicine, Eulji University, Daejeon 34824, Korea



Received: February 23, 2022
Revised: April 13, 2022
Accepted: May 14, 2022

[†]Corresponding author

Ki-Ho Lee
Dept. of Biochemistry and Molecular Biology, Eulji University, Daejeon 34824, Korea
Tel: +82-42-259-1643
Fax: +82-42-259-1649
E-mail: kiholee@eulji.ac.kr

Copyright © 2022 The Korean Society of Developmental Biology.
This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Yong-Seung Lee
<https://orcid.org/0000-0002-8442-4024>
Ki-Ho Lee
<https://orcid.org/0000-0002-3495-5126>

Conflict of interests

The authors declare no potential conflict of interest.

Acknowledgements

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2017R1D1A 1B03030209).

Authors' contributions

Conceptualization: Lee KH.
Data curation: Lee YS.
Formal analysis: Lee KH.
Methodology: Lee KH.
Investigation: Lee YS.

Abstract

The differentiation and development of preadipocyte into mature adipocyte are regulated by transcription factors, such as CCAAT enhancer binding protein (*Cebp*) gene family and sterol regulatory element binding transcription factor 1 (*Srebp1*). Steroid hormones give influences on the development and function of adipocyte. The present research examined expression patterns of CCAAT enhancer binding protein alpha (*Cebpa*), CCAAT enhancer binding protein beta (*Cebpb*), CCAAT enhancer binding protein gamma (*Cebpg*), sterol regulatory element binding transcription factor 1 (*Srebp1*), androgen receptor (*Ar*), and estrogen receptors (*Esr*) among different epididymal fat parts during postnatal period by quantitative real-time polymerase chain reaction. In the distal epididymal fat, expression of *Cebpa*, *Cebpb*, *Cebpg*, *Srebp1*, *Ar*, and *Esr2* was increased until 12 months of age, while expression of *Esr1* was decreased at 5 months of age and was not detectable after 8 months of age. In the proximal epididymal fat, transcript levels of *Cebps* and *Srebp1* were increased at 8 months of age, followed by decreases of *Cebpb* and *Cebpg* transcript levels at 12 months of age. An additional increase of *Srebp1* expression was observed at 12 months of age. Expression of *Ar* and *Esr2* were increased until 8 months of age, followed by a drop of *Ar* expression level at 12 months of age. Expression pattern of *Esr1* was similar to that in the distal epididymal fat. In the tail epididymal fat, expression of *Cebpa*, *Cebpg*, *Srebp1*, *Ar*, and *Esr2* was increased with age. *Esr1* was not detectable at all. The highest level of *Cebpb* was observed at 8 months of age. These data suggest the possibility of developmental and functional differentiation among the epididymal fat parts.

Keywords: Epididymal fat, Postnatal development, Real-time polymerase chain reaction (PCR), CCAAT enhancer binding protein, Sterol regulatory element binding transcription factor 1, Steroid hormone receptor

INTRODUCTION

The epididymal fat is a part of gonadal adipose tissue and is located nearby the epididymis, as named (Berry et al., 2013; Lee, 2020). Based on histochemical characteristics and/or relative localization, the

Writing-original draft: Lee KH.
Writing-review & editing: Lee KH, Lee YS.

Ethics approval

All procedures were carried out in accordance with the guidelines for the care and use of laboratory animals of National Research Council in Korea.

epididymal fat is discriminated into three types, distal, proximal, and tail epididymal fat (Tirard et al., 2007; Lee, 2020). The distal and proximal epididymal fat is the typically known epididymal fat pad, and differential gene expression between these epididymal fat parts has been observed from our previous and other researches, indicating functional differences between distal and proximal epididymal fat parts (Tirard et al., 2007; Lee & Kim, 2019; Lee, 2019). The tail epididymal fat is localized on the caudal epididymis and is continuously enlarged in size as aged (Lee, 2020). Even though the histochemical characteristics and function of tail epididymal fat have not been defined yet, expression patterns of some adipocyte-associated molecules in the tail epididymal fat are distinct from those detected in the distal and proximal epididymal fat pads (Lee, 2020). These observations suggest that the tail epididymal fat would have different functional roles from distal and proximal epididymal fat pads.

The epididymal fat is consisted of fully mature lipid-filled adipocyte and other types of cells, including preadipocyte, vascular cells, and extracellular matrix cells (Cleary et al., 1977). Like other fat tissues, the weight of epididymal fat is continuously increased during postnatal period (Cleary et al., 1977; Gruen et al., 1980), and such growth of epididymal fat is likely due to expansion of mature adipocyte in size and number (Cleary et al., 1977). Numerous factors, including transcription factors, micro RNAs, and growth factors, involve in development of adipocyte (Kuri-Harcuch et al., 2018). In addition, differentiation of preadipocyte into mature adipocyte and proliferation of preadipocyte are precisely controlled by a harmony of regulated expression of various adipogenic genes (Ntambi & Kim, 2000; Kuri-Harcuch et al., 2018). For examples, CCAAT enhancer binding protein (C/EBP) β seems to be the first transcription factor initiating the differentiation of preadipocyte into adipocyte and mediates the expression of C/EBP α at the later stage of differentiation (Ntambi & Kim, 2000). It is generally considered that C/EBP β also influences on the expression of peroxisome proliferator-activated receptor (PPAR) γ , which is a key regulator of downstream adipogenic gene expression for preadipocyte differentiation (Ntambi & Kim, 2000; Kuri-Harcuch et al., 2018). Even though differential expression patterns of adipocyte marker genes among different epididymal fat parts during postnatal development have been evaluated (Lee, 2019; Lee & Kim, 2019; Lee, 2020), the presence and expression patterns of differentiation-related genes of preadipocyte in the epididymal fat have not been examined in detail.

The effect of androgen and estrogen on the adipogenesis has also been extensively examined. It is generally considered that estrogen promotes hyperplasia of adipocyte and androgen likely involves in hypertrophy of adipocyte (Gavin & Bessesen, 2020). Because androgen and estrogen chiefly play their roles by binding with androgen receptor (AR) and estrogen receptor (ER), respectively, it is reasonable to examine expression of these receptors in the epididymal fat parts. In fact, Dieudonne et al. (1995) have demonstrated the presence of AR in the epididymal fat, especially higher level than in subcutaneous adipose tissue, and differential regulatory effect of testosterone via AR depending on proliferation and differentiation of preadipocytes. The expression of ER α (Metz et al., 2016) and ER β (Cheon et al., 2020) in the epididymal fat has also been shown. Together, even though a direct role of steroid hormones on the epididymal adipose tissue has not been revealed, expression of AR, ER α , and ER β in the epididymal fat pad suggests the possibility of functional role(s) of steroid hormones on the growth of epididymal fat pad during postnatal period.

Thus, the present research was focused to examine expression of transcription modulators involved in the differentiation of preadipocyte, such as C/EBP family genes [CCAAT enhancer binding protein alpha (*Cebpa*), CCAAT enhancer binding protein beta (*Cebpb*), and CCAAT enhancer binding protein gamma (*Cebpg*)] and sterol regulatory element binding transcription factor 1 (*Srebp1*), and steroid hormone receptors, including *Ar*, estrogen receptors (*Esr1*), and *Esr2*, in the different epididymal fat parts during postnatal period.

MATERIALS AND METHODS

1. Experimental animals and epididymal fat tissue collection

The C57BL/6N male mice at 1 month of postnatal age purchased from Samtako (OSan, Korea) were randomly separated and assigned into one of 4 postnatal age groups, 2 (n=6), 5 (n=6), 8 (n=6), and 12 (n=6) months of age. Free access to food and water of experimental animals under controlled environment were permitted for entire experimental period. All animal care procedures were performed in accordance with the guidelines for the care and use of laboratory animals of National Research Council in Korea.

The animal was euthanized by CO₂ stunning in a chamber at proper age, and the male reproductive tract was taken through a lower abdominal incision and was placed in cold PBS (Gibco, Grand Island, NY, USA). The epididymal fat pad separated from the epididymis was dissected and divided into the distal and proximal epididymal fats. The tail epididymal fat adjacent to caudal epididymis was collected and pooled in same age group to obtain total RNA extract sufficient for quantitative real-time polymerase chain reaction (PCR). Isolated epididymal fat tissues were quickly frozen in liquid nitrogen and stored in -80°C freezer for further use.

2. Isolation of total RNA and construction of complementary DNA (cDNA) strand

The total RNA pellet from epididymal fat tissue homogenized in TRI REAGENT solution (Molecular Research Center, Cincinnati, OH, USA) was extracted by consecutive exposure to chloroform and isopropanol. The total RNA was dissolved and resuspended in DNase/RNase-free dH₂O. The concentration of total RNA extract was determined by UV spectrophotometry (NanoDrop Lite, Thermo Scientific, Wilmington, DE, USA), and the general quality of total RNA extract was checked by 1.2% agarose gel electrophoresis.

One microgram of total RNA was utilized to generate the first-strand of cDNA with reverse transcription (RT) mixture (iScrip™ Reverse transcription Supermix for RT-qPCR, Bio-Rad Laboratories, Hercules, CA, USA) and nuclease free-dH₂O to adjust 20 µL in a final reaction volume. The condition of RT reaction was at 25°C for 5 min, 46°C for 20 min, and then 95°C for 1 min.

3. Performance of quantitative real-time PCR and statistical analysis of data

One microliter of cDNA solution was mixed with 10 pmol of each primer, 7 µL of iQ™ SYBR® Green Supermix (Bio-Rad Laboratories), and nuclease free-dH₂O to make a final PCR volume of 20 µL. The sequence and other information of each primer are summarized in Table 1. The PCR was carried out in a sequence of a pre-denaturation at 95°C for 5 min, cycles of a denaturation at 95°C for 30 sec, an annealing at T_m for 30 sec, and an extension at 72°C for 30 sec. An extension step at 72°C for 10 min was added at the end of each PCR. Agarose gel electrophoresis was always performed to check the size of PCR product. The 18s ribosomal RNA (*Rn18s*) gene was included as an internal control of PCR analysis.

Independently triplicated RT reactions and PCRs were accomplished to get a mean and standard error for each target gene at a particular postnatal age. Expression level of each molecule was normalized to that of *Rn18s*, and the data in a graph were presented in the relative ratio against transcript level of a specific gene at 2 months of age. The statistical comparison for expression level of a target gene among postnatal age groups was determined by one-way ANOVA. If there was a statistical significance present within experimental age groups for expression level of the target gene, Duncan's test, a post-hoc analysis, was then employed to evaluate statistical significance among different age groups. Less than 0.05 of *p*-value was considered as the existence of statistical

Table 1. Primer information for real-time PCR analysis

Gene	Sequence (5' → 3')	T _m (°C)	PCR product (bps)
<i>Cebpa</i> (NM_001287523.1)	(F) AGAAGTCGGTGGACAAGAAC (R) CTTGACCAAGGAGCTCTCAG	52	233
<i>Cebpb</i> (NM_009883.4)	(F) ATGCAATCCGGATCAAACGTG (R) CAACCCCGCAGGAACATCTTTA	60	113
<i>Cebpg</i> (NM_009884.3)	(F) ACCCATGGATCGGAATAGTG (R) GCTTCCAACCGTTCATTCTC	55	150
<i>Srebp1</i> (NM_011480.3)	(F) AGGTCACCGTTTCTTTGTGG (R) TCCAAGGGCATCTGAGAACT	55	208
<i>Ar</i> (NM_013476.4)	(F) TGTCAAAAGTGAAATGGGACC (R) TGGTACTGTCCAAACGCATGT	55	74
<i>Esr1</i> (AB560752.1)	(F) GCCTCTGGCTACCATTATGG (R) CATCTCTCTGACGCTTGTGC	55	244
<i>Esr2</i> (U81451.1)	(F) GTCAGGGTGGTTGGTTGTGG (R) ATCGCGTCACTTTCCTTTTCC	55	203
<i>Rn18s</i> (NM_03278.3)	(F) ACTCAACACGGGAAACCTCA (R) AACCAAGACAAATCGCTCCAC	55	123

PCR, polymerase chain reaction; *Cebpa*, CCAAT enhancer binding protein alpha; *Cebpb*, CCAAT enhancer binding protein beta; *Cebpg*, CCAAT enhancer binding protein gamma; *Srebp1*, sterol regulatory element binding transcription factor 1; *Ar*, androgen receptor; *Esr1*, estrogen receptor alpha; *Esr2*, estrogen receptor beta; *Rn18s*, 18s ribosomal RNA.

difference on the transcript level of the gene among different age groups, represented with different letters on bars in the graph.

RESULTS

1. Expression patterns of *Cebpa* and *Cebpb* in the distal, proximal, and tail epididymal fat during postnatal period

The expression level of *Cebpa* in the distal epididymal fat at 5 months of age was significantly higher than that at 2 months of age (Fig. 1A). A tremendous increase of *Cebpa* transcript level in the distal epididymal fat was observed at 8 months of age, followed by an additional increase of *Cebpa* expression at 12 months of age (Fig. 1A). In the proximal epididymal fat, expression level of *Cebpa* was not changed until 5 months of age (Fig. 1B). However, expression of *Cebpa* was significantly increased at 8 months of age, and there was no significant difference of *Cebpa* transcript level between 8 and 12 months of age (Fig. 1B). Like the proximal epididymal fat, there was no significant change of *Cebpa* expression at 5 months of age in the tail epididymal fat, compared with that at 2 months of age (Fig. 1C). An increase of *Cebpa* transcript level in the tail epididymal fat was detected at 8 months of age (Fig. 1C). Expression level of *Cebpa* in the tail epididymal fat at 12 months of age was significantly higher than that at 8 months of age (Fig. 1C).

Fig. 2 shows expression patterns of *Cebpb* in different epididymal fat parts at several postnatal ages. In the distal epididymal fat, expression level of *Cebpb* was significantly increased with age (Fig. 2A). But, there was no change of *Cebpb* transcript level in the proximal epididymal fat at 5 months of age, compared with the level at 2 months of age (Fig. 2B). Then, expression level of *Cebpb* in the proximal epididymal fat was significantly increased at 8 months of age, followed by a decrease at 12 months of age (Fig. 2B). Expression of *Cebpb* in the tail epididymal fat was significantly increased at 5 months of age, along with an additional increase of *Cebpb* transcript level at 8 months of age (Fig. 2C). However, there was no further significant change of *Cebpb* expression level at 12 months of age (Fig. 2C).




Fig. 1. Expression patterns of *Cebpa* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Cebpa* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Cebpa*, CCAAT enhancer binding protein alpha.




Fig. 2. Expression patterns of *Cebpb* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Cebpb* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Cebpb*, CCAAT enhancer binding protein beta.

2. Expression patterns of *Cebpg* and *Srebp1* in the distal, proximal, and tail epididymal fat during postnatal period

A significant increase of *Cebpg* in the distal epididymal fat was observed at 5 months of age, compared with at 2 months of age (Fig. 3A). Further increases of *Cebpg* transcript levels in the distal epididymal fat were also detected at 8 and 12 months of age (Fig. 3A). In the proximal epididymal fat, an increase of *Cebpg* expression was found at 8 months of age, followed by a significant drop at 12 months of age (Fig. 3B). Expression pattern of *Cebpg* in the tail epididymal fat was quietly similar with that in the distal epididymal fat, which showed continuous increase of *Cebpg* transcript level with age (Fig. 3C).

An increase of *Srebp1* transcript level in the distal epididymal fat was detected at 8 months of age (Fig. 4A). Expression level of *Srebp1* at 12 months of age was higher than that of 8 months of age (Fig. 4A). Expression pattern of *Srebp1* in the proximal epididymal fat was almost same with that in the distal epididymal fat (Fig. 4B). An increase of *Srebp1* transcript level was first observed at 8 months of age, followed by another significant surge of *Srebp1* expression at 12 months of age (Fig. 4B). The level of *Srebp1* transcript in the tail epididymal fat was significantly increased as aged (Fig. 4C).

3. Expression patterns of *Ar*, *Esr1*, and *Esr2* in the distal, proximal, and tail epididymal fat during postnatal period

Expression of *Ar* in the distal epididymal fat was significantly increased at 5 months of age




Fig. 3. Expression patterns of *Cebpg* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Cebpg* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Cebpg*, CCAAT enhancer binding protein gamma.




Fig. 4. Expression patterns of *Srebp1* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Srebp1* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Srebp1*, sterol regulatory element binding transcription factor 1.

(Fig. 5A). The level of *Ar* transcript in the distal epididymal fat was further increased at 8 months of age, followed by another increase of *Ar* transcript level at 12 months of age (Fig. 5A). The level of *Ar* transcript in the proximal epididymal fat at 5 months of age was also higher than 2 months of age (Fig. 5B). An additional increase of *Ar* expression level in the proximal epididymal fat was detected at 8 months of age, but a significant decrease of *Ar* transcript level at 12 months of age was followed in the proximal epididymal fat (Fig. 5B). Expression of *Ar* in the tail epididymal fat was significantly increased at 5 months of age, even though there was no change of *Ar* transcript level between 5 and 8 months of age (Fig. 5C). The abundance of *Ar* transcript in the tail epididymal fat at 12 months of age was higher than that at 8 months of age (Fig. 5C).

Expression of *Esr1* in the distal epididymal fat was first detected at 2 months of age, followed by a significant drop of *Esr1* transcript level at 5 months of age (Fig. 6A). The presence of *Esr1* expression in the distal epididymal fat at 8 and 12 months of age was not observed (Fig. 6A). Expression pattern of *Esr1* in the proximal epididymal fat during postnatal period was almost same as in the distal epididymal fat (Fig. 6B). There was no *Esr1* expression detected in the tail epididymal fat during the entire postnatal period (Fig. 6C).

Expression pattern of *Esr2* in different epididymal fat parts during postnatal period is shown in Fig. 7. Expression of *Esr2* in the distal epididymal fat was significantly increased at 5 months of age, followed by continuous increases at 8 and 12 months of age (Fig. 7A). In the proximal epididymal fat, the transcript level of *Esr2* at 5 months of age was higher than that at 2 months of age (Fig. 7B).




Fig. 5. Expression patterns of *Ar* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Ar* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Ar*, androgen receptor.




Fig. 6. Expression patterns of *Esr1* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Esr1* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; N.D., not detectable; *Esr1*, estrogen receptor alpha.




Fig. 7. Expression patterns of *Esr2* in the different epididymal fat parts of mouse at different postnatal ages. The relative transcript levels of *Esr2* in the distal (A), proximal (B), and tail (C) epididymal fat are shown. Different letters over solid bars indicate statistical significance at $p < 0.05$. M, months of postnatal age; *Esr2*, estrogen receptor beta.

A surge of *Esr2* transcript level was also detected at 8 months of age, followed by no change of *Esr2* expression level at 12 months of age (Fig. 7B). Expression of *Esr2* in the tail epididymal fat was also detected and significantly increased as aged (Fig. 7c).

DISCUSSION

The differentiation of preadipocyte into differentiated mature adipocyte requires the coordinated actions of a variety of transcription factors and modulators, as well as growth factors and hormones. It is generally acknowledged that the induction of C/EBP β and C/EBP δ in the early stage of adipogenesis promotes expression and activation of PPAR γ and C/EBP α , key regulators for establishing terminal differentiation of adipocyte and maintaining mature adipocyte phenotypes (Lee et al., 2018). In addition, SREBP1, a transcription factor involved in the differentiation of preadipocyte, participates in adipose differentiation via an increase of transcriptional activity of PPAR γ (Kim et al., 1998). Moreover, the concern of actions of androgen and estrogen on the development of epididymal fat tissue has been suggested (Dieudonne et al., 1995; Metz et al., 2016; Gavin & Bessesen, 2020). Thus, in the current research, expression patterns of these transcription factors and steroid hormone receptors within different epididymal fat regions during postnatal development have been examined using quantitative real-time PCR analysis.

The C/EBP family is a group of transcription factors involved in regulation of various physiological processes, including adipogenesis (Renfro et al., 2022). Of C/EBP proteins, C/EBP β is an initiation regulator for the differentiation of preadipocyte into mature adipocyte by stimulating expression of C/EBP α and PPAR γ (Lee et al., 2018). Then, C/EBP α and PPAR γ promote a variety of adipogenic factors, required for terminal differentiation of immature adipocyte into mature one and maintenance of mature status (Ali et al., 2013; Lee et al., 2018). Even though functional roles of C/EBP α and C/EBP β are extensively studied, less attention has been caught on molecular functions of C/EBP γ . It seems that C/EBP γ contributes its major roles on cell proliferation, development of hematopoietic and nervous system, and integrated cell stress response (Renfro et al., 2022). In addition, a possible role of C/EBP γ in energy homeostasis has been suggested (Renfro et al., 2022). Thus, it is supposed that C/EBP γ in adipocyte would involve in expression of marker molecules expressed from fully differentiated adipocyte and in conservation of mature status, rather than differentiation of preadipocyte. In the distal and tail epididymal fat parts, expression of *Cebpa*, *Cebpb*, and *Cebpg* is continuously increased until 12 months of age. These results have been expected because such expression patterns of *Cebp* genes would be necessary for increase of mature adipocyte in size and number with age (Cleary et al., 1977). However, expression of *Cebp* genes in the proximal epididymal fat between 2 and 5 months of age was not different. Moreover, transcript levels of *Cebp* genes at 12 months of age was not changed or even decreased, compared with that at 8 months of age. Such regional differences of the epididymal fat on expression patterns of *Cebp* genes during postnatal period imply that postnatal development of the proximal epididymal fat would be distinguishable from those of the distal and tail epididymal fat. In addition, to our knowledge, this is the first time to show expression patterns of *Cebpg* in the epididymal fat tissue. The physiological role of *Cebpg* in the epididymal fat is not clearly suggested at this point. Detailed molecular examination of *Cebp* genes in different epididymal fat parts is suggested to provide detailed information about functional roles of these genes.

Along with C/EBPs, SREBP1 is another factor acting in the early stage of differentiation of preadipocyte into adipocyte by stimulating expression of adipogenic genes (Kim & Spiegelman, 1996; Niemelä et al., 2008). Expression of *Srebp1* in the rat whole epididymal fat has been observed from our previous research, showing a higher level of *Srebp1* transcript level at the old age than that at the young age and the fluctuation of *Srebp1* expression during the early postnatal period (Lee & Kim, 2018). In the present research, the levels of *Srebp1* transcript in the distal and proximal epididymal fat remain unchanged, followed by significant increases at 8 and 12 months of age. On the other hand, expression of *Srebp1* in the tail epididymal fat was continuously increased during

postnatal period. Because the size and number of mature adipocyte become increased with age, it is reasonable to consider that extended and escalated action of SREBP1 in the epididymal fat would be required to express adequate amount of adipogenic genes at the old age. Examination of expression levels of adipogenic genes after selective inhibition of *Srebp1* expression would provide valuable information for functional role of SREBP1 in the epididymal fat.

The adipogenesis and development of adipose tissue are also regulated by testosterone and estrogen (Gavin & Bessesen, 2020). The effects of estrogen on the development of adipocyte are not straightforward. Estrogen promote proliferation of adipocyte precursor by inhibiting adipogenesis, while another evidence indicates stimulatory effect of estrogen on adipocyte differentiation (Newell-Fugate, 2017). Thus, Gavin & Bessesen (2020) have addressed that the effect of estrogen on adipocyte development is likely different, depending on the stage of adipocyte development. In spite of such complicate effects of estrogen on the development of adipocyte, it is generally agreed that estrogen exerts its effect via ESR1 (Newell-Fugate, 2017; Gavin & Bessesen, 2020). Testosterone via AR action has a suppressive effect on preadipocyte differentiation and overall reductive effect of adipose tissue mass (Newell-Fugate, 2017; Gavin & Bessesen, 2020). Increased expression of *Ar* and decreased expression of *Esr1* with age have been detected in the distal and proximal epididymal fat. Even no *Esr1* expression during postnatal period has been observed in the tail epididymal fat. An increase of *Esr2* transcription level with age has been found in all types of the epididymal fat. In fact, opposite expression patterns of *Ar* and *Esr1* in the epididymal fat have been expected. Such unexpected outcomes on *Ar* and *Esr1* expression patterns would be due to different types of adipose tissue. Indeed, differential responses depending on types of white adipose tissues (WAT) to testosterone and/or estrogen have been frequently suggested from other researches (Newell-Fugate, 2017; Gavin & Bessesen, 2020). Moreover, it is possible that these steroid hormones have unknown specific roles restricted to the development of epididymal fat. Thus, it is encouraged to examine the effects of testosterone and estrogen on postnatal development of the epididymal fat, which are separated from those on other types of adipose tissue.

The findings from the present study remain several questions to be answered for the development of epididymal fat during postnatal period. Despite, the current research has a value on the detection of differential expression patterns of several transcription factors-associated with preadipocyte differentiation and steroid hormone receptors in different epididymal fat parts at several postnatal ages. More attention has to be paid for evaluation of molecular biological roles of these transcription factors and steroid hormones within each epididymal fat part.

REFERENCES

- Ali AT, Hochfeld WE, Myburgh R, Pepper MS (2013) Adipocyte and adipogenesis. *Eur J Cell Biol* 92:229-236.
- Berry DC, Stenesen D, Zeve D, Graff JM (2013) The developmental origins of adipose tissue. *Development* 140:3939-3949.
- Cheon YP, Ko C, Lee KH (2020) Assessment of adipocyte differentiation and maturation-related gene expression in the epididymal fat of estrogen receptor α knockout (ER α KO) mouse during postnatal development period. *Dev Reprod* 24:287-296.
- Cleary MP, Greenwood MRC, Brasel JA (1977) A multifactor analysis of growth in the rat epididymal fat pad. *J Nutr* 107:1969-1974.
- Dieudonne MN, Pecquery R, Leneuve MC, Jaubert AM, Giudicelli Y (1995) Androgen receptors in cultured rat adipose precursor cells during proliferation and differentiation: Regional

- specificities and regulation by testosterone. *Endocrine* 3:537-541.
- Gavin KM, Bessesen DH (2020) Sex differences in adipose tissue function. *Endocrinol Metab Clin North Am* 49:215-228.
- Gruen R, Kava R, Greenwood MR (1980) Development of basal lipolysis and fat cell size in the epididymal fat pad of normal rats. *Metabolism* 29:246-253.
- Kim JB, Spiegelman BM (1996) ADD1/SREBP1 promotes adipocyte differentiation and gene expression linked to fatty acid metabolism. *Genes Dev* 10:1096-1107.
- Kim JB, Wright HM, Wright M, Spiegelman BM (1998) ADD1/SREBP1 activates PPAR γ through the production of endogenous ligand. *Proc Natl Acad Sci USA* 95:4333-4337.
- Kuri-Harcuch W, Velez-delValle C, Vazquez-Sandoval A, Hernández-Mosqueira C, Fernandez-Sanchez V (2019) A cellular perspective of adipogenesis transcriptional regulation. *J Cell Physiol* 234:1111-1129.
- Lee JE, Schmidt H, Lai B, Ge K (2018) Transcriptional and epigenomic regulation of adipogenesis. *Mol Cell Biol* 39:e00601-e00618.
- Lee KH, Kim NH (2018) Expressional patterns of adipocyte-associated molecules in the rat epididymal fat during postnatal development period. *Dev Reprod* 22:351-360.
- Lee KH, Kim NH (2019) Differential expression of adipocyte-related molecules in the distal epididymal fat of mouse during postnatal period. *Dev Reprod* 23:213-221.
- Lee KH (2019) Postnatal expressional patterns of adipose-associated molecules in the mouse proximal epididymal fat. *Dev Reprod* 23:313-322.
- Lee KH (2020) Expression of adipocyte-associated genes in the mouse tail epididymal fat at different postnatal ages. *Dev Reprod* 24:167-176.
- Metz L, Gerbaix M, Masgrau A, Guillet C, Walrand S, Boisseau N, Boirie Y, Courteix D (2016) Nutritional and exercise interventions variably affect estrogen receptor expression in the adipose tissue of male rats. *Nutr Res* 36:280-289.
- Newell-Fugate AE (2017) The role of sex steroids in white adipose tissue adipocyte function. *Reproduction*. 153:R133-R149.
- Niemälä S, Miettinen S, Sarkanen JR, Ashammakhi N (2008) Adipose tissue and adipocyte differentiation: Molecular and cellular aspects and tissue engineering applications. In: Ashammakhi N, Reis R, Chiellini F (eds), *Topics in Tissue Engineering*, Vol. 4, Biomaterials and Tissue Engineering Group, Oulu, Finland, pp 1-26.
- Ntambi JM, Kim YC (2000) Adipocyte differentiation and gene expression. *J Nutr* 130:3122S-3126S.
- Renfro Z, White BE, Stephens KE (2022) CCAAT enhancer binding protein gamma (C/EBP- γ): An understudied transcription factor. *Adv Biol Regul* 84:100861.
- Tirard J, Gout J, Lefrançois-Martinez AM, Martinez A, Begeot M, Naville D (2007) A novel inhibitory protein in adipose tissue, the aldo-keto reductase AKR1B7: Its role in adipogenesis. *Endocrinology* 148:1996-2005.